


What is claimed is:

1. A compound having the structure:

5

10



15

wherein X is CR₇; N; or N⁺O⁻;

wherein Y is O; CO; S; CR₃R₅; or NR₆;

20

wherein each R₂ is independently H; F; Cl; Br; I; -NO₂, -CN; straight chained or branched C₁-C₄ alkyl; C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy; -OH; -(CH₂)_qOH; -COR₄; CO₂R₄; CONHR₄; phenyl; or benzyl;

25

wherein each R₃ is independently H; straight chained or branched C₁-C₄ alkyl; C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy; -(CH₂)_qOH; -OH; =N-OR₄; COR₄; CO₂R₄; CONHR₄; phenyl; or benzyl;

30

wherein each R₄ is independently H; straight chained or branched C₁-C₄ alkyl, C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; or phenyl;

wherein each R_5 is independently H; straight chained or branched C_1 - C_4 alkyl, C_1 - C_4 monofluoroalkyl, or C_1 - C_4 polyfluoroalkyl;

5 wherein R_6 is H; straight chained or branched C_1 - C_4 alkyl; C_1 - C_4 monofluoroalkyl or C_1 - C_4 polyfluoroalkyl; straight chained or branched C_1 - C_4 alkoxy; -
 $CH_2CH_2(CH_2)_qOH$; COR_4 ; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

10 wherein each R_7 is independently H; -CN; straight chained or branched C_1 - C_4 alkyl; C_1 - C_4 monofluoroalkyl or C_1 - C_4 polyfluoroalkyl; straight chained or branched C_1 - C_4 alkoxy; -OH; -
 $(CH_2)_qOH$; - COR_4 ; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

15 wherein m and n are each independently 0, 1, 2 or 3, provided that m+n is 2 or 3;


20 wherein each p is independently 0, 1 or 2; and

wherein each q is independently 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

25

2. A compound having the structure:



wherein each of Z1, Z2 and Z3 is N or CR₂, with the proviso that either one of Z1, Z2 or Z3 is N and the others of Z1, Z2 or Z3 are CR₂, or both Z1 and Z3 are N and Z2 is CR₂;

wherein R₁ is H; F; straight chained or branched C₁-C₄ alkyl, C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy, -OH; or -(CH₂)_qOH;

wherein each R₂ is independently H; F; Cl; Br; I; -NO₂, -CN; straight chained or branched C₁-C₄ alkyl; C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy; -OH; -(CH₂)_qOH; -COR₄; CO₂R₄; CONHR₄; phenyl; or benzyl;

wherein each R₄ is independently H; straight chained or branched C₁-C₄ alkyl, C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; or phenyl; and


wherein q is each independently 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.


3. The compound of claim 1 or 2, wherein the compound comprises the (+) enantiomer.
- 5 4. The compound of claim 1 or 2, wherein the compound comprises the (-) enantiomer.
5. The compound of claim 1, wherein Y is CR_3R_5 , and m+n is 3.
- 10 6. The compound of claim 1, wherein Y is CR_3R_5 and m+n is 2.
7. The compound of claim 1, wherein Y is NR_6 .
- 15 8. The compound of claim 1, wherein X is N.
9. The compound of claim 2, wherein two of Z1, Z2 and Z3 are CR_2 and the other is N.
- 20 10. The compound of claim 5, wherein p is at least 1 and at least one R_3 is methyl.
11. The compound of claim 5, wherein at least one R_2 is methyl.
- 25 12. The compound of claim 6, wherein at least one R_2 is bromo.
13. The compound of any one of claims 10, 11, or 12, wherein X is N.
- 30 14. The compound of claim 9, wherein at least one R_2 is methyl or phenyl.

- 5

- 10




- 20



30

18. The compound of claim 6 having the structure:


5



10

19. The compound of claim 12 having the structure:

15




20

20. The compound of claim 15 having the structure:

25

30




21. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 1 or 2 and a pharmaceutically acceptable carrier.

5 22. A method for treating an α_2 adrenergic receptor associated disorder in a subject, which comprises administering to the subject an amount of a compound effective to treat the disorder, wherein the compound has the structure:

10

15



20

wherein X is CR_7 ; N ; or N^+O^- ;

wherein Y is O ; CO ; S ; CR_5R_5 ; or NR_5 ;

25

wherein each R_2 is independently H ; F ; Cl ; Br ; I ; $-NO_2$; $-CN$; straight chained or branched C_1 - C_4 alkyl; C_1 - C_4 monofluoroalkyl or C_1 - C_4 polyfluoroalkyl; straight chained or branched C_1 - C_4 alkoxy; $-OH$; $-(CH_2)_4OH$; $-COR_4$; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

30

wherein each R_3 is independently H ; straight chained or branched C_1 - C_4 alkyl; C_1 - C_4 monofluoroalkyl or C_1 - C_4 polyfluoroalkyl; straight chained or branched C_1 - C_4 alkoxy; $-(CH_2)_4OH$; $-OH$; $=N-OR_4$; COR_4 ; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

35

wherein each R_4 is independently H; straight chained or branched C_1-C_4 alkyl; C_1-C_4 monofluoroalkyl or C_1-C_4 polyfluoroalkyl; or phenyl;

5 wherein each R_5 is independently H; straight chained or branched C_1-C_4 alkyl, C_1-C_4 monofluoroalkyl, or C_1-C_4 polyfluoroalkyl;

10 wherein R_6 is H; straight chained or branched C_1-C_4 alkyl; C_1-C_4 monofluoroalkyl or C_1-C_4 polyfluoroalkyl; straight chained or branched C_1-C_4 alkoxy; -
 $CH_2CH_2(CH_2)_qOH$; COR_4 ; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

15 wherein each R_7 is independently H; -CN; straight chained or branched C_1-C_4 alkyl; C_1-C_4 monofluoroalkyl or C_1-C_4 polyfluoroalkyl; straight chained or branched C_1-C_4 alkoxy; -OH; $-(CH_2)_qOH$; $-COR_4$; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

20 wherein m and n are each independently 0, 1, 2 or 3, provided that m+n is 2 or 3;


wherein each p is independently 0, 1 or 2; and

25 wherein each q is independently 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

30

23. A method for treating an α_2 adrenergic receptor associated disorder in a subject, which comprises administering to the subject an amount of a compound effective to treat the disorder, wherein the compound has the structure:



wherein each of Z1, Z2 and Z3 is N or CR₂, with the proviso that either one of Z1, Z2 or Z3 is N and the others of Z1, Z2 or Z3 are CR₂, or both Z1 and Z3 are N and Z2 is CR₂;

wherein R₁ is H; F; straight chained or branched C₁-C₄ alkyl, C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy, -OH; or -(CH₂)_qOH;

wherein each R₂ is independently H; F; Cl; Br; I; -NO₂, -CN; straight chained or branched C₁-C₄ alkyl; C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy; -OH; -(CH₂)_qOH; -COR₄; CO₂R₄; CONHR₄; phenyl; or benzyl;


wherein each R₄ is independently H; straight chained or branched C₁-C₄ alkyl, C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; or phenyl; and

wherein q is each independently 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

5 24. The method of claim 22 or 23, wherein the disorder is migraine headache, hypertension or glaucoma.

25. A method for treating pain in a subject, which comprises administering to the subject an amount of a compound effective to treat the subject's pain, wherein
10 the compound has the structure:



wherein X is CR₇; N; or N⁺O⁻;

wherein Y is O; CO; S; CR₃R₅; or NR₅;

25 wherein each R₂ is independently H; F; Cl; Br; I; -NO₂, -CN; straight chained or branched C₁-C₄ alkyl; C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy; -OH; -(CH₂)₃OH; -COR₄; CO₂R₄; CONHR₄; phenyl; or benzyl;
30

wherein each R₃ is independently H; straight chained or branched C₁-C₄ alkyl; C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄

alkoxy; $-(CH_2)_qOH$; $-OH$; $=N-OR_4$; COR_4 ; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

5 wherein each R_4 is independently H; straight chained or branched C_1-C_4 alkyl, C_1-C_4 monofluoroalkyl or C_1-C_4 polyfluoroalkyl; or phenyl;

10 wherein each R_5 is independently H; straight chained or branched C_1-C_4 alkyl, C_1-C_4 monofluoroalkyl, or C_1-C_4 polyfluoroalkyl;

15 wherein R_6 is H; straight chained or branched C_1-C_4 alkyl; C_1-C_4 monofluoroalkyl or C_1-C_4 polyfluoroalkyl; straight chained or branched C_1-C_4 alkoxy; $-CH_2CH_2(CH_2)_qOH$; COR_4 ; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

20 wherein each R_7 is independently H; $-CN$; straight chained or branched C_1-C_4 alkyl; C_1-C_4 monofluoroalkyl or C_1-C_4 polyfluoroalkyl; straight chained or branched C_1-C_4 alkoxy; $-OH$; $-(CH_2)_qOH$; $-COR_4$; CO_2R_4 ; $CONHR_4$; phenyl; or benzyl;

25 wherein m and n are each independently 0, 1, 2 or 3, provided that $m+n$ is 2 or 3;


wherein each p is independently 0, 1 or 2; and

wherein each q is independently 0, 1, 2 or 3;

30 or a pharmaceutically acceptable salt thereof.

26. A method for treating pain in a subject, which comprises administering to the subject an amount of a compound effective to treat the subject's pain, wherein the compound has the structure:

5



10

wherein each of Z1, Z2 and Z3 is N or CR₂, with the proviso that either one of Z1, Z2 or Z3 is N and the others of Z1, Z2 or Z3 are CR₂, or both Z1 and Z3 are N and Z2 is CR₂;

15

wherein R₁ is H; F; straight chained or branched C₁-C₄ alkyl, C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy, -OH; or - (CH₂)_qOH;

20

wherein each R₂ is independently H; F; Cl; Br; I; -NO₂, -CN; straight chained or branched C₁-C₄ alkyl; C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; straight chained or branched C₁-C₄ alkoxy; -OH; - (CH₂)₃OH; -COR₄; CO₂R₄; CONHR₄; phenyl; or benzyl;

25

wherein each R₄ is independently H; straight chained or branched C₁-C₄ alkyl, C₁-C₄ monofluoroalkyl or C₁-C₄ polyfluoroalkyl; or phenyl; and

30

wherein q is each independently 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

35